


BIOLOGY, LEGAL STATUS, CONTROL MATERIALS, AND DIRECTIONS FOR USE

House Mice

Mus musculus

Family: Muridae



Introduction: House mice are not native to North America but are classed as an invasive species imported on early settler and trade ships. Worldwide house mice are some of the most troublesome and damaging rodents. In Australia they are a preoccupation of the agricultural community costing millions of dollars in damage. Population explosions can see as many as 1000 mice per hectare. The West coast of the United States is no exception, although, as yet, not to the same epic proportions seen in Australia. House mice transmit parasites and diseases to other animals

(including people), consume and contaminate food, and damage structures and property as a result of their gnawing activities. They are resilient and thrive under a wide variety of climates and conditions. They are well adapted to living in close contact with humans and thrive in the conditions that man provides, particularly where hygiene and housekeeping standards are low. House mice are much more common in residences and commercial structures than rats (Brooks 1973). It is important that mouse infestations are controlled.



Identification: House mice are small and agile rodents with a slightly pointed nose, and small black protruding eyes. They are usually a uniform grey/ grayish brown color, and sometimes the lower half of their bodies may be a slightly lighter shade. They have slender bodies, 5 to 7 inches long, and weigh ½ to 1 ounce. They have rather large sparsely haired ears for their body size, and a semi hairless tail which is

about as long as the entire body and head.

House mice nibble on many foodstuffs, discard partially eaten items and create a mess with droppings, urine, and hairs. They will nest build. Outside house mice living in fields may dig up and feed on newly planted grain, or cause damage by eating crops before harvest. Because of the similarities with deer mice and meadow voles; correct identification before embarking on any pest control plan will ensure efficient control.

Visual sightings of mice are of limited value in accurately estimating the population present. Search premises thoroughly when looking for mice, in structures, attics, basements, around foundations, crawl spaces, and behind and under stored materials.

It may be possible to detect house mice presence using non toxic tracking dust or alternatively flour or talcum powder at 20 – 30 foot intervals throughout a structure (Salmon, Whisson, Marsh, 2006)



Legal Status: House mice are classified as nongame mammals by the California Fish and Game Code. House Mice threatening or injuring growing crops or other property of which you are the owner or tenant, you may control them using any legal means. They may also be taken by Department of Food and Agriculture, federal state, and county officers or employees, when acting in their official capacities.




Damage: House mice will contaminate the environments in which they live. House mice are omnivorous; they consume and contaminate stored grain and cereal products as well as fruit, vegetables, cheese, meat, and hay. They are often a pest in poultry facilities where they consume the chicken feed, and can damage electrical equipment.

House mice carry a wide range of diseases which can be passed on to humans. House mice and their ectoparasites are involved in the transmission of diseases: salmonella food poisoning by infected rodent feces on suitable foods; rickettsialpox by the bite of the house mouse mite; and lymphocytic choriomeningitis, a virus infection of house mice, may be transmitted to man by means of food or dust contaminated with respiratory droplets or powdered feces of infected animals.

House mice will inevitably cause physical property damage whenever present in and about buildings. This may be costly and potentially dangerous creating fire or safety hazards. The damage is seen in the form nest building and gnawing of structures, electrical wire insulation, rubber insulation, stored items in basements, attics, garages i.e. paintings, books, clothing, furniture, family heirlooms.

House mice damage can be identified by referencing: feces/droppings, gnawing, markings or smudges



against solid structure (wood beams), nests, odor, sounds, track, visual sighting.



Range: House mice are found throughout the entire United States and are considered the most common mammal in cities, except for humans (Brooks 1973).

House Mice



Habitat: Occasionally found in fields, but usually in buildings. House mice build nests in boxes and crates as well as stationary objects, enhancing their rapid distribution along transportation routes.



Biology: House mice live in a wide variety of man-made structures as well as in open fields. Mice living in the open may move into buildings when weather becomes severe. House mice soon become accustomed to ordinary noises and, as a result, may be active in areas near people, animals, or machines.

House mice have poor vision but a keen sense of smell, taste, hearing and touch. The long whiskers on the nose and above the eyes serve the sense of touch. Mice normally make their runways close to walls, using their whiskers as feelers. House mice eat the same foods as man, showing a preference for foods high in protein or carbohydrates. Different environments offer varied diets and sometimes mice will feed upon such things as live insects, starch in clothing, and glue in bookbinding's. They require very little water and can live for many months on a dry grain diet. Because of their small size, they must feed several times during a 24-hour period. This means they will be active during the day as well as the night. Their range is normally 10 to 30 feet from the nest. The nest is lined with soft material such as cotton or paper and may be built in walls, cabinets, upholstered furniture or other convenient space. The usual length of life is about one year. During this time, a female house mouse will raise an average of 30 to 35 young; sexual maturity is attained at 1-1/2 months; gestation period is 18-21 days; and several litters (up to 8 per year) of 3 to 11, usually 5 or 6, young are raised. The young are born blind.



Damage Prevention and Control Methods: Controlling house mice differs substantially from controlling rats. Essentially house mice are smaller, have a smaller home range, reproduce faster than rats, and in many cases are less susceptible to rodenticides. Any control program should integrate three components: sanitation measures and habitat modification, structure proofing and exclusion, and population control.

Exclusion: Physical barriers can prevent mice from gaining entry to structures where food and shelter are available. "Rodent proofing" is an important and often neglected aspect of rodent control. It is a relatively permanent form of control which can prevent damage from occurring.

To exclude mice, seal all holes and openings larger than 1/4-inch across. Rodent proofing should be done with heavy materials that will resist rodent gnawing. These include concrete mortar, galvanized sheet metal, and heavy-gauge hardware cloth.

Habitat Modification and Sanitation: Sanitation, which includes good housekeeping practices, proper storage and handling of food, feed and garbage, is stressed as a method of rodent control. Unfortunately, even the best sanitation will not eliminate house mice. It will, however, aid in control by permitting easier detection of mouse sign, increasing effectiveness of traps and baits by reducing competing food items, and by preventing mice from flourishing and reaching high populations.

Although house mice are less dependent upon humans for their existence than are Norway rats, they are much more adaptable to living with people. They require very little space and only small amounts of food to exist. Mice have been known to inhabit buildings even before construction has been complete, living off the crumbs and scraps of worker lunches. In offices, mice may live behind cabinets or furniture and feed on scraps or crumbs from lunches and snacks and on food kept in desks. In homes, they may find ample food in kitchens, and in the garage, they will eat sacked or spilled pet food, grass seed, or insects such as cockroaches. Thus, no matter how good the sanitation, most buildings in which food is stored, prepared, or consumed will support at least a few mice. For this reason, a constant watch must be kept for mice which may invade the premises.



Where possible, store bulk foods in rodent-proof containers or rooms. Stack boxed foods in orderly rows on pallets in a way that allows for thorough inspection for evidence of mice. In such storage areas, keep stored materials away from walls. A 12-inch white band painted on the floor next to the wall serves as a reminder to keep items away from walls. It will allow you to detect rodent droppings or

other signs easily. Sweep floors frequently to permit ready detection of fresh droppings.

When storing food or feed on pallets, keep in mind that mice can jump up more than 12 inches from a flat surface. They are also good climbers and can walk up surfaces such as wood or concrete (unless they have a slick finish). Mice can live for a considerable period of time within a pallet of feed and only rarely come down to the floor.

Regular removal of debris and control of weeds from around structures will reduce the amount of shelter available to rodents. In some instances, a strip of heavy gravel placed adjacent to building foundations or other structures will reduce rodent burrowing in these locations. In any event, keeping the periphery of buildings and other structures clean of weeds and debris (including stacked lumber, firewood, and other stored materials) will discourage rodent activity and will allow easier detection of sign.

Frightening: This is not considered a practicable solution for house mouse control. Various commercial ultrasonic devices claim success at controlling house mice but no scientific research exists to support their effectiveness in controlling mice. What is known about ultrasound is that it is very directional and therefore does not travel well around corners; it also loses its intensity rapidly as it leaves its source. Testing has shown that house mice may be driven away for a few days but return and resume normal activity.

Repellents: House mice like all rodents find some types of taste and odors objectionable. However, chemical repellants are not a practicable solution for mouse infestations. Moth balls (naphthalene) or household ammonia (bleach) may have a short term effect although they are not registered as repellents with the EPA. A number of repellents are available which claim to prevent gnawing by mice or other rodents but their effectiveness is questionable.

Trapping: Trapping can be an effective method of controlling mice, but it requires more labor than most other methods. Trapping is recommended where poisons are unadvisable or when mice need to be removed from inhabited buildings. It is the preferred method to try first in homes, garages, and other small structures where there may be only a few mice present.

Trapping has several advantages: 1) it does not rely on rodenticides; 2) it permits the user to view the results; and 3) it allows for disposal of the trapped mice, thereby eliminating odor problems from decomposing carcasses which may occur when poisoning is done within buildings.

The simple, inexpensive, wood based snap trap is available in most hardware and farm supply stores. Traps should be baited with a small piece of nutmeat, chocolate candy, dried fruit, or bacon, tied or glued securely to the trigger. Peanut butter or marshmallows also may be used as bait. Food baits that become stale lose their effectiveness. Set traps close to walls, behind objects, in dark places, and in locations where mouse activity is seen. Place the traps so that when mice follow their natural course of travel (usually close to a wall) they will pass directly over the trigger. Set traps so that the trigger is sensitive and will spring easily. When traps are set in runways or in travel routes, effectiveness can be increased by enlarging the trigger. This can be done with a square of cardboard, metal or screen wire that fits just inside the wire deadfall. Some traps are already equipped with this type of trigger.

Use enough traps to make the campaign short and decisive. Mice seldom venture far from their shelter and food supply, so traps should be spaced no more than about six feet apart in areas where mice are active. Although mice are not nearly as afraid of new objects as rats, leaving the traps baited but unset until the bait is taken at least once will reduce the chance of mice escaping the trap and becoming trap-shy.

A variety of different mouse traps designed as lethal traps and also non lethal are available in most retail stores. The efficacy of these traps is dependent largely upon their placement. Use as many traps as practicable so that trapping time will be short. As many as a dozen or more traps may be necessary for a heavily infested home or other building structure. Mice rarely travel far from their shelter and food supply.

Dispose of dead mice by burying or placing them in plastic bags in the trash. When handling any rodent carcass always wear waterproof gloves to protect yourself from disease. Do not touch them with bare hands and wash thoroughly after handling traps.

Glue boards: An alternative to traps are glue boards, which catch and hold mice attempting to cross them much the same way flypaper catches flies. Place glue boards wherever mice travel -- along walls or in

established pathways. Do not use glue boards where children, pets, or wildlife can contact them. Glue boards lose their effectiveness in dusty areas unless covered, and temperature extremes may affect the tackiness of some glue. They are considered less effective in capturing rats than mice. You can purchase ready-to-use glue boards, or you can buy glue to make your own boards or traps. Dispose of live trapped rodents in a humane manner.

REFERENCES AND ADDITIONAL READING

Advani, Ranjan, 1992. Field Evaluation of Three Anticoagulant Rodenticides Against *Mus musculus* Populations in Apartment Buildings in New York City. Proc. 15th Vertebrate Pest Conf. (J.E. Borrecco & R. E. Marsh, Eds.) Published at Univ. of Calif., Davis. Pp. 208-211.

Brown, Peter R., G.R. Singleton, D.A. Jones, S.C. Dunn, 1998. The Management of House Mice in Agricultural Landscapes Using Farm Management Practices: An Australian Perspective. Proc. 18th Vertebrate Pest Conf. (R.O. Baker & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 156-159.

Caughley, Judy, C. Donkin, K.Strong, 1998. Managing Mouse Plagues in Rural Australia. Proc. 18th Vertebrate Pest Conf. (R.O. Baker & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 160-165.

Colvin, Bruce A., A.D. Ashton, W.C. McCartney, W.B. Jackson, 1990. Planning Rodent Control For Boston's Central Artery/Tunnel Project. Proc.14th Vertebrate Pest Conf. (L.R. Davis and R.E. Marsh, Eds.) Published at Univ. of Calif., Davis Pp. 65-69.

Corrigan, Robert M., C.A. Towell, R.E. Williams, 1992. Development of Rodent Control Technology for Confined Swine Facilities. Proc. 15th Vertebrate Pest Conf. (J.E. Borrecco & R. E. Marsh, Eds.) Published at Univ. of Calif., Davis. Pp. 280-285.

Corrigan, Robert M., 1998. The Efficacy of Glue Traps Against Wild Populations Of House Mice, *Mus Domesticus*, Ruddy. Proc. 18th Vertebrate Pest Conf. (R.O. Baker & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 268-275.

Corrigan, Robert M., 2004. An Overview of the Significance and Management of Vertebrate Pests Around Zoological Parks. Proc. 21st Vertebrate Pest Conf. (R.M. Timm and W.P. Gorenzel, Eds.) Published at Univ. of Calif., Davis. Pp. 327-337.

Corrigan, Robert M., D.C. Collins, 2004. The Possible Effects of Bait Container Design on Mouse Feeding Activity in Real-World Structural Baiting Situations. Proc. 21st Vertebrate Pest Conf. (R.M. Timm and W.P. Gorenzel, Eds.) Published at Univ. of Calif., Davis. Pp. 174-179.

Doane, Becky, D. Blodget, B. Bonnivier, 1996. How To Control A Pest's Pest-Flea and Rodent Efficacy. Proc. 17th Vertebrate Pest Conf. (R.M. Timm & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 197-198.

Frantz, Stephen C., C. Padula Madigan, 1998. Warfarin Resistance Revisited. Proc. 18th Vertebrate Pest Conf. (R.O. Baker & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 276-280.

Hinds, Lyn A., 2006. Immunocontraception of Small Mammals: Case Study for the Wild House Mouse in Australia. Proc. 22nd Vertebrate Pest Conf. (R.M. Timm and J.M. O'Brien, Eds.) Published at Univ. of Calif., Davis. Pp. 96-101.

Hosea, Robert C., 2000. Exposure of Non-Target Wildlife to Anticoagulant Rodenticides in California. Proc. 19th Vertebrate Pest Conf. (T.P. Salmon & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 236-244.

Humphries, Richard E., A.P. Meehan, R.M. Sibly, 1992. The Characteristics and History of Behavioral Resistance in Inner-City House Mice (*Mus domesticus*) in the U.K. Proc. 15th Vertebrate Pest Conf. (J.E. Borrecco & R. E. Marsh, Eds.) Published at Univ. of Calif., Davis. Pp. 161-164.

Hygnstrom, Scott E., 1992. Impacts of House Mouse Activity on Five Types of Insulation. Proc. 15th Vertebrate Pest Conf. (J.E. Borrecco & R. E. Marsh, Eds.) Published at Univ. of Calif., Davis. Pp. 286-288.

Hygnstrom, Scott E., K.C. VerCauteren, R.M. Timm, B.M. Corrigan, J.G. Beller, L.L. Bitney, M.C. Brumm, D. Meyer, D.R. Virchow, R.W. Wills, 2002. An Economic Model of Integrated House Mouse Control in Swine production Facilities. Proc. 20th Vertebrate Pest Conf. (R.M. Timm and R.H. Schmidt, Eds.) Published at Univ. of Calif., Davis. Pp. 53-58.

Jacobs, William W., 1990. Required Use of Protective Bait Stations in the U.S. Proc. 14th Vertebrate Pest Conf. (L.R. Davis and R.E. Marsh, Eds.) Published at Univ. of Calif., Davis. Pp. 36-42.

Kaukeinen, D.E., C.W. Spragins, J.F. Hobson, 2000. Risk-Benefit Considerations in Evaluating Commensal Anticoagulant Rodenticide Impacts to Wildlife. Proc. 19th Vertebrate Pest Conf. (T.P. Salmon & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 245-256.

- Marshall, Edward F., 1992. The Effectiveness of Difethialone (LM 2219) for Controlling Norway Rats And House Mice Under Field Conditions. Proc. 15th Vertebrate Pest Conf. (J.E. Borrecco & R. E. Marsh, Eds.) Published at Univ. of Calif., Davis. Pp. 171-174.
- Mach, Jeff J., 2004. Investigations of Commensal Rodenticide Bait against Wild Norway Rats Plus Additional Toxicology Data of Warfarin on Laboratory Norway Rats and House Mice Proc. 21st Vertebrate Pest Conf. (R.M. Timm and W.P. Gorenzel, Eds.) Published at Univ. of Calif., Davis. Pp. 140-144.
- McCann, Geraldine R., 2000. Chlorophacinone and Diphacinone: Standard *Mus Musculus* and *Peromyscus Maniculatus* Anticoagulant Laboratory Tests. Proc. 19th Vertebrate Pest Conf. (T.P. Salmon & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 263-267.
- Miller Baker, Ann Eileen, 1994. Stowaway Transport Rates of House Mice (*Mus domesticus*) and Deermice (*Peromyscus maniculatus*). Proc. 16th Vertebrate Pest Conf. (W.S. Halverson & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 106-112.
- O'Brien, John M., R.E. Marsh, 1990. Vertebrate Pests of Beekeeping. Proc. 14th Vertebrate Pest Conf. (L.R. Davis and R.E. Marsh, Eds.) Published at Univ. of Calif., Davis Pp. 228-232.
- Prescott, Colin V., 1996. Preliminary Study of the Genetics of Resistance in the House Mouse. Proc. 17th Vertebrate Pest Conf. (R.M. Timm & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 83-87.
- Silberhorn, Eric M., J.F. Hobson, G.H. Miller, N.J. Condos, 2000. U.S. EPA Reregistration Eligibility Decision (Red) for the Rodenticide Cluster: Overview of the Regulatory Process, Response of Registrants and Stakeholders, and Implications for Agricultural and Urban Rodent Control. Proc. 20th Vertebrate Pest Conf. (R.M. Timm and R.H. Schmidt, Eds.) Published at Univ. of Calif., Davis. Pp. 268-276.
- Singleton, Grant R., 2000. Ecologically-Based Rodent Management Integrating New Developments in Biotechnology. Proc. 19th Vertebrate Pest Conf. (T.P. Salmon & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 221-227.
- Sorensen, Ronald B., R.C. Nuti, M.C. Lamb, 2006. Rodent Management for Surface Drip Irrigation Tubing in Peanut. Proc. 22nd Vertebrate Pest Conf. (R.M. Timm and J.M. O'Brien, Eds.) Published at Univ. of Calif., Davis. Pp. 201-205.
- Timm, Robert M., 1983. Prevention and Control of Wildlife Damage. University of Nebraska Cooperative Extension Service. pp. B32-B42.
- Tobin, Mark E., R.T. Sugihara, R.M. Engeman, 1994. Effects of Initial Rat Capture on Subsequent Capture Success Of Traps. Proc. 16th Vertebrate Pest Conf. (W.S. Halverson & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 101-105.
- Von Wahlde, Matt, B.A. Colvin, 1994. Using Geographical Information Systems for Tracking an Urban Rodent Control Program. Proc. 16th Vertebrate Pest Conf. (W.S. Halverson & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 327-334.

Witmer, Gary W., H. Martins, L. Flor, 2004. Leptospirosis in the Azores: the Rodent Connection. Proc. 21st Vertebrate Pest Conf. (R.M. Timm and W.P. Gorenzel, Eds.) Published at Univ. of Calif., Davis. Pp. 217-220.

Witmer, Gary, S. Jojola, 2006. What's Up with House Mice? – A Review. Proc. 22nd Vertebrate Pest Conf. (R.M. Timm and J.M. O'Brien, Eds.) Published at Univ. of Calif., Davis. Pp. 124-130.