


*BIOLOGY, LEGAL STATUS, CONTROL MATERIALS, AND DIRECTIONS FOR USE*

**Roof Rat**

*Rattus rattus*, **Black Rat**

Family: Muridae



**Introduction:** The roof rat, like the Norway rat, is an introduced species in the United States. A third rat species, the Polynesian rat (*R. exulans*) is present in the Hawaiian Islands but not on the mainland United States. The Roof rat is commonly known as *Rattus rattus*, the black rat, and ship rat. Roof rats were common on early sailing ships which accounts for their distribution in the USA. Roof rats have a long history as carriers of plague.



**Identification:** Roof Rats (*Rattus rattus*), sometimes called black rats, are slightly smaller than Norway rats. Unlike Norway rats, their tails are longer than their heads and bodies combined. Roof rats are very agile climbers and usually live and nest above ground in shrubs, trees, and dense vegetation such as ivy. In buildings, they are most often found in enclosed or elevated spaces in attics, walls, false ceilings, and cabinets. The roof rat has a more limited geographical range than the Norway rat, preferring ocean-influenced, warmer climates. In areas where the roof rat occurs, the

Norway rat may also be present. If you are unsure of the species, look for rats at night with a strong flashlight or trap a few. There are several key physical differences between the two species of rats. See Table 1.

**Table 1. Identifying Characteristics of Adult Rats.**

Characteristic	Roof rat	Norway rat
general appearance	sleek, agile	Large, robust
color of belly	gray to white	mostly grayish
body weight	5 to 10 ounces	7 to 18 ounces
tail	extends at least to snout; black, fine scales	shorter than body; dark above; pale below; scales
head	muzzle pointed	muzzle blunt
ears	long enough to reach eyes   folded over	do not reach eyes



**Legal Status:** Roof rats are classified as nongame mammals by the California Fish and Game Code. Nongame mammals which are found to be injuring growing crops or other property may be taken at any time or in any manner by the owner or tenant of the premises. They may also be taken by officers or employees of the department of Food and Agriculture or by federal or county officers or employees when acting in their official capacities pursuant to the provisions of the Food and Agricultural Code pertaining to pests.



**Damage:** Essentially the same as the Norway rat (see discussion in preceding section) from economic and public health standpoints.

In some agricultural areas, roof rats cause significant damage and loss to tree crops e.g. citrus, avocados, walnuts, almonds, and other nuts. They often eat all the pulp from oranges while the fruit is still hanging from the tree, leaving only the empty rind. With lemons they may eat the rind and leave only the fruit. They may eat the bark and girdle smaller branches. In sugarcane fields they move into fields and eat the stalks. Roof rats may also eat seeds, or seedlings of plants. Vegetable, fruit, melon berry can also suffer minor damage.



Like the Norway rat the roof rat can transmit diseases to humans including murine typhus, leptospirosis, salmonellosis (food poisoning), rat bite fever, and plague. It is also capable of transmitting diseases to domestic animals and is implicated in the transference of ectoparasites as well.



**Range:** Range: The "roof rat" whose coloration resembles that of the Norway rat, lives along the coast, in the interior valleys and in mountains to 5,000 feet. The

"black rat" is a color variant of this species. It is found mainly near salt water. Some sources separate the "roof rat" into two additional subdivisions, the Alexandrine rat and the fruit rat, based on characteristic coloration differences.

### Black Rat



**Habitat:** Buildings, especially upper stories and attics, ships, warehouses, fields, stream banks, dense vines, palms and other trees.



**Biology:** Roof rats are smaller than Norway rats, and the tail is usually longer than the head and body combined. The roof rat is less aggressive and has been displaced by the Norway rat in some parts of the United States. The roof rat is a more agile climber than his larger cousin and it seems better able to establish in some rural habitats. The roof rat is more likely to choose enclosed spaces in attics, walls, and cabinets for nesting and cover. The roof rat is omnivorous but shows more preference for fruits and vegetables than does the Norway rat; both like cereals. The roof rat's home range is generally 100 to 150 feet.

The reproductive potential of roof rats is almost as high as that of Norway rats; an average of about 20 young per year are weaned by each female. Litter size averages six to eight young and an average of four to six litters are raised per year. The gestation period is about 22 days and the young may run about at three weeks of age. Sexual maturity is reached at three to five months and the average length of life is about one year. Like the Norway rat, much of the population dies out and is replaced in less than a year. Both rats breed year round.



### Damage Prevention and Control Methods

**Rat Damage Indicators:** Droppings can be found along 'runways' in feeding areas, and near shelter. They may be as large as  $\frac{3}{4}$  inch long, and  $\frac{1}{4}$  inch wide. Fresh droppings are soft in texture.

Tracks can include footprints or tail marks, and can often be seen on dusty surfaces or mud. Rats can be tracked by placing a small amount of flour in a patch across a suspected runway area overnight. The presence of tracks indicates rodents.

Urine both wet and dry will fluoresce under ultraviolet light. Urine stains may occur along travelways or in feeding areas.

Runs or burrows can be found next to walls, along fences, next to buildings, under bushes or debris. Rats memorize pathways and use the same routes habitually.

Smudge marks (rub marks) can occur on beams, rafters, pipes, and walls due to oil and dirt rubbing off a rat's fur as it frequently travels routes.

Gnawing is a basic rat activity. One way rats keep their paired incisor teeth worn down is by gnawing on hard surfaces. Size of entry holes differ from mice, rat entry holes are often 2 inches whereas mice are slightly smaller.

Rats can often be heard climbing in walls, gnawing, clawing, squeaking, and making fighting noises.

**Exclusion:** Physical barriers are an excellent way to prevent rats from gaining entry to structures where food and shelter are available.

Rats can be excluded by sealing all holes and openings larger than  $\frac{1}{2}$  inch. Use strong materials that will resist rodent gnawing; concrete mortar, galvanized sheet metal, steel wool, and heavy-gauge hardware cloth.

To prevent rodents from climbing or traveling along a particular route, install guards made of sheet metal or similar materials. Guards must be wide enough and positioned to keep rodents from reaching their outer margins by climbing or jumping.

Sheet metal band attached to a wall will prevent climbing by rodents. Rodent guards should be at least 12 inches to 18 inches wide. Inside buildings, such guards can prevent rats from climbing at corners. Guards also can be installed to prevent rodents from climbing the outside of buildings having rough exterior walls. When used in combination with hardware cloth or other suitable material, they can make a building essentially rodent-proof. This technique has been used to make corn cribs, barns, and other older buildings rat-proof.

**Habitat Modification:** Good sanitation is an excellent way to minimize any rat pest problems. The elimination of food and water through good warehouse sanitation can do much towards reducing rodent infestation. Proper garbage and refuse disposal containers along with an exterior sanitation program are also very helpful. Emphasis should be placed on the removal of as much harborage as is practical.

Dense shrubbery, vine-covered trees and fences, and vine ground cover all make ideal harborage for roof rats. Pruning and/or removal of certain ornamentals are often required to obtain a degree of lasting rat

control.

Poor sanitation is one of the basic reasons for the continued existence of moderate to high rat populations in urban and suburban areas. In agricultural environments, proper sanitation cannot always eliminate rat populations, but it can often prevent rats from flourishing in large numbers.

Sanitation involves good housekeeping, including proper storage and handling of food materials, feed and edible garbage. Warehouses, granaries and grain mills, silos, port facilities, and similar structures may provide excellent habitat for rats. Store bulk foods in rodent-proof containers or rooms. Stack sacked or boxed foods in orderly rows on pallets in a way that allows thorough inspection for evidence of rats. In such storage areas, keep stored materials away from walls. A 12-inch white band painted on the floor adjacent to the wall will aid in detecting rodent droppings and other rat sign. Sweep floors frequently to permit ready detection of fresh rat presence..

Pet foods often are a source of food for rats in and around homes. Keep all such materials stored and subsequently removed for disposal. Proper refuse storage containers are rat-and damage-resistant, and equipped with a tight-fitting lid. Racks or stands prevent corrosion or rusting of containers, reduce rat shelter under containers, and minimize the chance of containers being overturned.



Bulk storage containers for refuse, such as those used at apartments, businesses, and housing projects, should be similarly rodent-proof. Large metal refuse containers (dumpsters) sometimes have drain holes to facilitate cleaning. These drain holes should be fitted with a wire

mesh screen or a removable plug; otherwise, the container becomes a large feeding station for rodents.

**Frightening:** Naturally rats are wary and frighten easily when they encounter unfamiliar sounds. However, most rodents, including rats, rapidly become accustomed to new sounds when heard repeatedly. Temporary success may be achieved, and rats may be repelled from an immediate area, but they will ultimately return and resume their normal activities.

Many commercially available devices produce ultrasonic sound to frighten, claiming it controls rodents. Research shows that rodents may be repelled temporarily from an immediate area, but will return and resume normal activities in the same way any new sound will affect the rodents. Ultrasonic devices are often expensive and their effectiveness is questionable. They are not recommended as a solution to rodent problems.

### **Non Anticoagulants:**

There are four non-anticoagulant rodenticides registered by the EPA for control of roof rats: bromethalin, cholecalciferol (vitamin D<sub>3</sub>), red squill, and zinc phosphide. However, only zinc phosphide is registered for agricultural use. All can be used for controlling anticoagulant-resistant populations of rats. Where rat numbers are large costs of baiting with non anticoagulants may reduce overall costs.

Bromethalin (Assault®, Vengeance®) is formulated in ready-to-use bait as a chronic rodenticide, applied so that rats will have the opportunity to feed on the bait one or more times over a period of one to several days. Because it is a slow-acting compared to zinc phosphide or red squill, bait shyness is not usually a problem, nor is prebaiting necessary to get good control in most situations.

Cholecalciferol (vitamin D<sub>3</sub>, Quintox®) is similarly formulated in ready-to-use bait, serving as a chronic rodenticide. Death occurs 3 or 4 days after ingestion of a lethal dose. Because the toxicant is slow-acting, bait shyness is not reported to occur. It is claimed that rodents cease feeding once a lethal dose has been ingested.

Red squill is a selective and relatively safe toxicant for use only against Norway rats. It acts as an emetic, which provides some degree of protection to certain nontarget species that might accidentally consume the bait. Rats, which cannot vomit, are unable to rid themselves of the toxicant once it is consumed. In the past, one problem was the variation in the quality of the material, which is derived from a plant. Red squill must be stored in a sealed container, as moisture will cause it to loss potency. Use for more than a few days at a time may result in bait shyness.

Zinc phosphide is a dark gray powder, insoluble, that has been used extensively in the control of rodents. It is available in ready-to-use dry baits and also in concentrates to prepare fresh baits. Its strong garlic-like odor appears to be attractive to rodents that are not bait-shy.

**Baits:** Baits are produced as pellets, treated grain or meal. Sugar, to five percent by weight, is sometimes added to improve bait acceptance by both roof rats and Norway rats.

**Trapping:** Trapping can be an effective method of controlling rats, but it requires more skill and labor than most other methods. Trapping is recommended where poisons are inadvisable. It is the preferred method to try first in homes, garages, and other small structures where there may be only a few rats present.



Trapping has several advantages: 1) it does not rely on inherently hazardous rodenticides; 2) it permits the user to view success; and 3) it allows for disposal of the rat carcasses, thereby eliminating odor problems from decomposing carcasses which may remain when poisoning is done within buildings.

The simple, inexpensive, wood-based snap trap is available in most

hardware and farm supply stores. Traps should be baited with a small piece of hot dog, bacon, or nutmeat tied securely to the trigger. Peanut butter or marshmallows also may be used as bait. Baits that become stale lose their effectiveness.

Set traps close to walls, behind objects, in dark corners, and in places where roof rat activity is seen. This will usually be in attic areas, along fences and possibly in trees. Place the traps so that when rats follow their natural course of travel (usually close to a wall) they will pass directly over the trigger. Set traps so that the trigger is sensitive and will spring easily. When traps can be set in runways or in travel routes, effectiveness can be increased by enlarging the trigger. This can be done with a square of cardboard, metal, or screen wire that fits just inside the wire deadfall. Some commercial traps come with enlarged triggers. If trapping outside, take care so that birds and small animals cannot enter the traps.

Use enough traps to make the campaign short and decisive. Leaving traps unset until the bait has been taken at least once reduces the chance of rats escaping the trap and becoming trap-shy.

Other kinds of traps are also effective in catching rats. Wire-mesh cage traps such as the Tomahawk® and Havahart® can be used effectively to capture rats alive. Wire funnel-entrance traps have also been used for this purpose. These live catch traps do present the problem of disposing of the trapped animal. Fish and Game regulations prevent the release of live animals without a permit. Euthanize with CO<sub>2</sub>.

Keep traps clean and in good working condition. When dirty, clean them in a hot detergent solution with a stiff brush. Human and dead-rat odors on traps are not known to reduce trapping success.

An alternative to traps are glue boards, which catch and hold rats attempting to cross them much the same way flypaper catches flies. Place glue boards wherever rats travel -- along walls, or in established pathways. Do not use glue boards where children, pets, or wildlife can contact them. Glue boards lose their effectiveness in dusty areas unless covered, and temperature extremes may affect the tackiness of some glue. They are considered less effective for capturing rats than for mice. You can purchase ready-to-use glue boards, or you can buy glue to make your own boards or traps. Dispose of live trapped rodents in a humane manner; euthanize live, trapped rodents by asphyxiation with carbon dioxide, or use a stick to kill them with a sharp blow to the base of the skull.

**Directions for Use** (with reference primarily to farms and other agricultural or field use)

Spot Baiting (except anticoagulant baits): Follow label instructions. Generally, place a teaspoon quantity of bait (in each active burrow or scatter small amounts of bait in protected places frequented by rats, but inaccessible to livestock, poultry, wildlife and children. Whenever practical, prebaiting several days before applying acute or one-shot toxic bait will achieve better control and will give an indication of how much toxic bait to put out. Prebaiting should always be conducted where natural food is abundant.

Bait should be picked up and disposed of upon completion of rodent control program. Do not retreat with zinc phosphide baits for at least three months.

**Tracking Powder:** Toxic dusts or powders have been used for many years to control rats and mice. When rats walk over a patch of toxic powder, they pick some of it up on their feet and fur and later ingest it while grooming. Tracking powders are useful in controlling rats where food is plentiful and good bait acceptance is difficult to achieve. Rats are more likely to ingest a lethal amount of a poorly-accepted toxicant applied by this method than if it is mixed into a bait material. There is little likelihood of toxicant shyness developing when using tracking powders.

## REFERENCES AND ADDITIONAL READING

- Buckle, A.P., M.G.P. Fenn, 1992. Rodent Control in the Conservation of Endangered Species. Proc. 15th Vertebrate Pest Conf. (J.E. Borrecco & R. E. Marsh, Eds.) Published at Univ. of Calif., Davis. Pp. 36-41.
- Buckle, A.P., C.V. Prescott, K.J. Ward, 1994. Resistance to the First and Second Generation Anticoagulant Rodenticides—A New Perspective. Proc. 16th Vertebrate Pest Conf. (W.S. Halverson & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp.138-144.
- Colvin, Bruce A., R. Degregorio, C Fleetwood, 1996. Norway Rat Infestation of Urban Landscaping and Preventive Design Criteria. Proc. 17th Vertebrate Pest Conf. (R.M. Timm & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 165-171.
- Colvin, Bruce A., 2000. A Comprehensive Rodent Control Program for Washington, D.C. Proc. 20th Vertebrate Pest Conf. (R.M. Timm and R.H. Schmidt, Eds.) Published at Univ. of Calif., Davis. Pp. 339-345.
- Corrigan, Robert M., 2004. An Overview of the Significance and Management of Vertebrate Pests around Zoological Parks. Proc. 21st Vertebrate Pest Conf. (R. M. Timm and W. P. Gorenzel, Eds.) Published at Univ. of Calif., Davis. Pp. 327-337.
- Childs, James E., T.G. Ksiazek, P.E. Rollin, J.W. Krebs, S.Zari, S.T. Nichol, C.J. Peters, G.E. Glass, 1994. Hantaviruses and Their Rodent Reservoirs in the United States. Proc. 16th Vertebrate Pest Conf. (W.S. Halverson & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 188-191.
- Doane, Becky, D. Blodget, B. Bonnivier, 1996. How to Control A Pest's Pest—Flea and Rodent Efficacy. Proc. 17th Vertebrate Pest Conf. (R.M. Timm & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 197-198.

Frantz, Stephen C., C. Padula Madigan, 1998. Warfarin Resistance Revisited. Proc. 18th Vertebrate Pest Conf. (R.O. Baker & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 276-280.

Hadler, Malcolm R., A.P. Buckle, 1992. Forty Five Years of Anticoagulant Rodenticides – Past, Present and Future Trends. Proc. 15th Vertebrate Pest Conf. (J.E. Borrecco & R. E. Marsh, Eds.) Published at Univ. of Calif., Davis. Pp. 149-155.

Hueth, Brent, D.Cohen, D. Zilberman, 1998. Non-Predator Vertebrate Pest Damage in California Agriculture: An Assessment of Economic Impacts in Selected Crops. Proc. 18th Vertebrate Pest Conf. (R.O. Baker & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 371-377.

Jacobs, William W., 1990. Required Use of Protective Bait Stations in the U.S. Proc.14th Vertebrate Pest Conf. (L.R. Davis and R.E. Marsh, Eds.) Published at Univ. of Calif., Davis Pp. 36-42.

Kaukeinen, D.E., C.W. Spragins, J.F. Hobson, 2000. Risk-Benefit Consideration in Evaluating Commensal Anticoagulant Rodenticides Impacts to Wildlife. Proc. 20th Vertebrate Pest Conf. (R.M. Timm and R.H. Schmidt, Eds.) Published at Univ. of Calif., Davis. Pp. 245-256.

Key, Gillian, K. Hudson, 2000. The Rat Control Program on the Island of St. Helena. Proc. 20th Vertebrate Pest Conf. (R.M. Timm and R.H. Schmidt, Eds.) Published at Univ. of Calif., Davis. Pp. 133-138.

Miller, Lowell A., K.A. Fagerstone, 2000. Induced Infertility As A Wildlife Management Tool. Proc. 20th Vertebrate Pest Conf. (R.M. Timm and R.H. Schmidt, Eds.) Published at Univ. of Calif., Davis. Pp. 160-168.

Miller, Lowell A., J. Rhyan, G. Killian, 2004. GonaCon, a Versatile GnRH Contraceptive for a Large Variety of Pest Animal Problems. Proc. 21st Vertebrate Pest Conf. (R. M. Timm and W. P. Gorenzel, Eds.) Published at Univ. of Calif., Davis. Pp. 269-273.

Morgan, D.R., G.R. Wright, S.C. Ogilvie, 1996. Assessment of the Environmental Impact of Brodifacoum During Rodent Eradication Operations in New Zealand. Proc. 17th Vertebrate Pest Conf. (R.M. Timm & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 213-218.

Prescott, C.V., M. El-Amin, R.H. Smith, 1992. Calciferols and bait Shyness in the Laboratory Rat. Proc. 15th Vertebrate Pest Conf. (J.E. Borrecco & R. E. Marsh, Eds.) Published at Univ. of Calif., Davis. Pp. 218-223.

Silberhorn, Eric M., J.F. Hobson, G.H. Miller, N.J. Condos, 2000. U.S. EPA Reregistration Eligibility Decision (Red) for the Rodenticide Cluster: Overview of the Regulatory Process, Response of Registrants and Stakeholders, and Implications for Agricultural and Urban Rodent Control. Proc. 20th Vertebrate Pest Conf. (R.M. Timm and R.H. Schmidt, Eds.) Published at Univ. of Calif., Davis. Pp. 268-276.

Witmer,, Gary W., E.W. Campbell III, F.Boyd, 1998. Rat Management For Endangered Species Protected in the U.S. Virgin Islands. Proc. 18th Vertebrate Pest Conf. (R.O. Baker & A.C. Crabb, Eds.) Published at Univ. of Calif., Davis. Pp. 281-286.

Witmer, Gary W., H. Martins, L. Flor, 2004. Leptospirosis in the Azores: the Rodent Connection. Proc. 21st Vertebrate Pest Conf. (R.M. Timm and W.P. Gorenzel, Eds.) Published at Univ. of Calif., Davis. Pp. 217-220.