Oral Dosage Forms That Should Not Be Crushed

John F. Mitchell, PharmD, FASHP¹

Last updated: October 2012

| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|---------------------------------|-----------------------------------|----------------|--|
| AcipHex | RABEprazole | Tablet | Extended-release |
| Actiq | fenta NYL | Lozenge | Slow-release Note: this lollipop delivery system requires the patient to slowly dissolve in mouth |
| Actonel | risedronate | Tablet | Irritant Note: chewed, crushed, or sucked tablets may cause oropharyngeal irritation |
| Adalat CC | NIFEdipine | Tablet | Extended-release |
| Adderall XR | amphetamine salts | Capsule | Extended-release (a) |
| AeroHist Plus | combination | Tablet | Slow-release (h) |
| Afeditab CR | NIFEdipine | Tablet | Extended-release |
| Afinitor | everolimus | Tablet | Mucous membrane irritant |
| Aggrenox | combination | Capsule | Extended-release |
| Alavert Allergy (Sinus 12 Hour) | combination | Tablet | Extended-release |
| Allegra-D | combination | Tablet | Extended-release |
| ALPRAZolam ER | ALPRAZ olam | Tablet | Extended-release |
| Altoprev | lovastatin | Tablet | Extended-release |
| Ambien CR | zolipidem | Tablet | Extended-release |
| Amibid DM | combination | Tablet | Extended-release |
| Amitiza | lubiprostone | Capsule | Slow-release |
| Ampyra | dalfampridine | Tablet | Extended-release Note: formerly fampridine-SR |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|------------------------|-----------------------------------|----------------|--|
| Amrix | cyclobenzaprine | Capsule | Extended-release |
| Aplenzin | bu PROP ion | Tablet | Extended-release |
| Apriso | mesalamine | Capsule | Extended-release (a) |
| | | | Note: maintain pH at less than or equal to 6.0 |
| Aptivus | tipranavir | Capsule | Note: oil emulsion within spheres; taste |
| Aquatab C | combination | Tablet | Slow-release (h) |
| Aquatab D | combination | Tablet | Slow-release (b) |
| Aricept 23 mg | donepezil | Tablet | Note: crushing the 25 mg tablet may significantly increase the rate of absorption; the 5 and 10 mg tablet are not affected |
| Arthrotec | diclofenac | Tablet | Delayed-release; Enteric-coated |
| Asacol | mesalamine | Tablet | Slow-release |
| Aspirin enteric-coated | aspirin | Tablet, Caplet | Delayed-release; Enteric-coated |
| Atelvia | risedronate | Tablet | Extended-release; Note: tablet coating is an important part of the delayed release |
| Azulfidine EN | sulfa SALA zine | Tablet | Delayed-release |
| Augmentin XR | combination | Tablet | Extended-release (b,h) |
| AVINza | morphine | Capsule | Enteric-coated (a; not pudding) |
| Avodart | dutasteride | Capsule | Note: drug may cause fetal abnormalities; women who are, or may become, pregnant, should not handle capsules; all women should use caution in handling capsules, especially leaking capsules |
| Bayer Regular | aspirin | Caplet | Enteric-coated |
| Biaxin-XL | clarithromycin | Tablet | Extended-release |
| Bidex A | combination | Tablet | Extended-release |
| Bidhist-D | combination | Tablet | Extended-release |
| Biltricide | praziquantel | Tablet | Taste (h) |
| Biohist LA | combination | Tablet | Extended-release (h) |
| Bisa-Lax | combination | Tablet | Enteric-coated (c) |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|--------------------|-----------------------------------|-----------------|---|
| Bisac-Evac | bisacodyl | Tablet | Enteric-coated (c) |
| Bisacodyl | combination | Tablet | Enteric-coated (c) |
| Boniva | ibandronate | Tablet | Note: chewed, crushed, or sucked tablets may |
| | | | cause oropharyngeal irritation |
| Bromfed PD | combination | Capsule | Extended-release (b) |
| Budeprion SR | combination | Tablet | Extended-release |
| Calan SR | verapamil | Tablet | Extended-release (h) |
| Carbatrol | car BAM azepine | Capsule | Extended-release (a) |
| Cardene SR | ni CAR dipine | Capsule | Extended-release |
| Cardizem | diltiazem | Tablet | Note: although not in the PI, the drug has a |
| | | | coating that is intended to release the drug over |
| | | | approximately 3 hours |
| Cardizem CD | diltiazem | Capsule | Extended-release |
| Cardizem LA | diltiazem | Tablet | Extended-release |
| Cardura XL | doxazosin | Tablet | Extended-release |
| Cartia XT | diltiazem | Capsule | Extended-release |
| Cefaclor ER | combination | Tablet | Extended-release |
| Ceftin | cefuroxime | Tablet | Taste (b) |
| | | | Note: use suspension for children |
| Cefuroxime | combination | Tablet | Taste (b) |
| | | | Note: use suspension for children |
| CellCept | mycophenolate | Capsule, Tablet | Teratogenic potential (i) |
| Charcoal Plus | charcoal, activated | Tablet | Enteric-coated |
| Chlor-Trimeton | combination | Tablet | Extended-release (b) |
| Cipro XR | ciprofloxacin | Tablet | Extended-release (b) |
| Claritin-D | combination | Tablet | Extended-release (b) |
| Claritin-D 24 Hour | combination | Tablet | Extended-release |
| Colace | docusate | Capsule | Taste (e) |
| Colestid | colestipol | Tablet | Slow-release |
| Concerta | methylphenidate | Tablet | Extended-release |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|--------------------|-----------------------------------|----------------|--|
| Commit | nicotine | Lozenge | Note: integrity compromised by chewing or crushing |
| Cotazym-S | pancrelipase | Capsule | Enteric-coated (a) |
| Covera-HS | verapamil | Tablet | Extended-release |
| Creon | pancrelipase | Capsule | Extended-release (a) |
| Crixivan | indinavir | Capsule | Note: capsule may be opened and mixed with fruit puree (e.g., banana) |
| Cymbalta | DULoxetine | Capsule | Extended-release (a) Note: may add contents of capsule to apple juice or applesauce but NOT chocolate |
| Cytoxan | cyclophosphamide | Tablet | Note: drug may be crushed but company recommends using injection |
| Cytovene | ganciclovir | Capsule | Skin irritant |
| Depakene | divalproex | Capsule | Slow-release; mucous membrane irritant (b) |
| Depakote | divalproex | Tablet | Delayed-release |
| Depakote ER | divalproex | Tablet | Extended-release |
| Depakote Sprinkles | divalproex | Capsule | Extended-release (a) |
| Detrol LA | tolterodine | Capsule | Extended-release |
| Dexilant | dexlansoprazole | Capsule | Delayed-release (a) |
| Dilacor XR | diltiazem | Capsule | Extended-release |
| Dilatrate-SR | isosorbide | Capsule | Extended-release |
| Dilt-CD | diltiazem | Capsule | Extended-release |
| Diltia XT | diltiazem | Capsule | Extended-release |
| Ditropan XL | oxybutynin | Tablet | Extended-release |
| Divalproex ER | combination | Tablet | Extended-release |
| Doxidan | bisacodyl | Tablet | Enteric-coated (c) |
| Drisdol | ergocalciferol | Capsule | Liquid filled (d) |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|------------------------------|-----------------------------------|-----------------|--|
| Droxia | hydroxyurea | Capsule | Note: exposure to the powder may cause serious skin toxicities; healthcare workers should wear gloves to administer |
| Dulcolax | bisacodyl | Tablet; Capsule | Enteric-coated (c); Liquid-filled |
| DynaCirc CR | isradipine | Tablet | Extended-release |
| EC-Naprosyn | combination | Tablet | Delayed-release; enteric-coated |
| Ecotrin (all) | aspirin | Tablet | Enteric-coated |
| E.E.S. 400 | erythromycin | Tablet | Enteric-coated (b) |
| Effer-K | potassium bicarbonate | Tablet | Effervescent tablet (f) |
| Effervescent Potassium | | Tablet | Effervescent tablet (f) |
| Effexor XR | venlafaxine | Capsule | Extended-release |
| Embeda | morphine sulfate | Capsule | Extended-release (a); do not give via N/G tube |
| E-Mycin | erythromycin | Tablet | Enteric-coated |
| Enablex | darifenacin | Tablet | Slow-release |
| Entocort EC | budesonide | Capsule | Extended-release; Enteric-coated (a) |
| Equetro | car BAM azepine | Capsule | Extended-release (a) |
| Ergomar | ergotamine | Tablet | Sublingual form (g) |
| Erivedge | vismodegib | Capsule | Note: PI indicates potential teratogenic; MSDS warns against skin contact; healthcare workers should take appropriate cautions |
| Ery-Tab | erythromycin | Tablet | Delayed-release; Enteric-coated |
| Erythromycin Stearate | | Tablet | Enteric-coated |
| Erythromycin Base | | Tablet | Enteric-coated |
| Erythromycin Delayed-Release | | Capsule | Enteric-coated Pellets (a) |
| Evista | raloxifene | Tablet | Taste; teratogenic potential (i) |
| Exalgo | HYDRO morphone | Tablet | Extended-release Note: breaking, chewing, crushing, dissolving before swallowing or injecting could result in overdose |
| Extendryl (all) | combination | Capsule | Extended-release (b) |
| Feen-a-mint | bisacodyl | Tablet | Enteric-coated (c) |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|--------------------------------------|-----------------------------------|----------------|--|
| Feldene | piroxicam | Capsule | Mucous membrane irritant |
| Fenta NYL | | Lozenge | Slow-release |
| | | | Note: this lollipop delivery system requires the |
| | | | patient to slowly dissolve in mouth |
| Fentora | fenta NYL | Tablet | Note: buccal tablet; swallow whole |
| Feosol | ferrous sulfate | Tablet | Enteric-coated (b) |
| Feratab | ferrous sulfate | Tablet | Enteric-coated (b) |
| Fergon | ferrous gluconate | Tablet | Enteric-coated |
| Fero-Grad 500 mg | combination | Tablet | Slow-release |
| Ferro-Sequels | combination | Tablet | Slow-release |
| Flagyl ER | metro NIDAZOLE | Tablet | Extended-release. |
| Fleet Laxative | bisacodyl | Tablet | Enteric-coated (c) |
| Flomax | tamsulosin | Capsule | Slow-release |
| Focalin XR | dexmethylphenidate | Capsule | Extended-release (a) |
| Fosamax | alendronate | Tablet | Mucous membrane irritant |
| Gleevec | imatinib | Tablet | Taste (h) |
| | | | Note: may be dissolved in water or apple juice |
| Glucophage XR | met FORMIN | Tablet | Extended-release |
| Glucotrol XL | glipi ZIDE | Tablet | Extended-release |
| Glumetza | met FORMIN | Tablet | Extended-release |
| Gralise | gabapentin | Tablet | Extended-release |
| Guaifed | combination | Capsule | Extended-release |
| Guaifed-PD | combination | Capsule | Extended-release |
| Guai FEN esin/Pseudoephedrine | | Tablet | Extended-release |
| Guaifenex DM | combination | Tablet | Extended-release (h) |
| Guaifenex GP | combination | Tablet | Extended-release |
| Guaifenex PSE | combination | Tablet | Extended-release (h) |
| Guaimax-D | combination | Tablet | Extended-release |
| Halfprin 81 | aspirin | Tablet | Enteric-coated |
| Horizant | gabapentin | Tablet | Extended-release |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|---------------------------------|-----------------------------------|---------------------|---|
| Hista-Vent DA | combination | Tablet | Extended-release (h) |
| Hydrea | hydroxyurea | Capsule | Note: exposure to the powder may cause serious skin toxicities; healthcare workers should wear gloves to administer |
| Imdur | isosorbide | Tablet | Extended-release (h) |
| Inderal LA | propranolol | Capsule | Extended-release |
| Indomethacin SR | indomethacin | Capsule | Extended-release (a,b) |
| InnoPran XL | propranolol | Capsule | Extended-release |
| Intelence | etravirine | Tablet | Note : tablet should be swallowed whole and not crushed; tablet may be dispersed in water |
| Intermezzo | zolpidem | Tablet (sublingual) | (g) |
| Intuniv | guan FACINE | Tablet | Extended-release |
| Invega | paliperidone | Tablet | Extended-release |
| Isoptin SR | verapamil | Tablet | Extended-release (h) |
| Isordil Sublingual | isosorbide | Tablet (sublingual) | (g) |
| Isosorbide Dinitrate Sublingual | isosorbide | Tablet (sublingual) | (g) |
| Isosorbide SR | isosorbide | Tablet | Extended-release |
| ISO tretinoin | ISO tretinoin | Capsule | Mucous membrane irritant |
| Jakafi | ruxolitinib | Tablet | Note: see PI for making a suspension |
| Jalyn | dutasteride / tamsulosin | Capsule | Note: women who are, or may become, pregnant, should not handle crushed or broken tablets (I) |
| Janumet XR | sitaGLIPtin/ metFORMIN | Tablet | Extended-release |
| Kadian | morphine | Capsule | (a) Extended-release Note: may add contents of capsule to applesauce without crushing) |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|----------------------|-----------------------------------|----------------|--|
| Kaletra | lopinavir/ritoavir | Tablet | Film-coated (b) |
| | | | Note: active ingredients are surrounded by a wax |
| | | | matrix to prevent healthcare exposure; women |
| | | | who are, or may become, pregnant, should not |
| | | | handle crushed or broken tablets |
| Kapidex | dexlansoprazole | Capsule | Delayed-release (a) |
| Kapvay | clo NID ine | Tablet | Extended-release |
| Keppra | lev ETIRA cetam | Tablet | Taste (b) |
| Keppra XL | lev ETIRA cetam | Tablet | Extended-release (b) |
| Ketek | telithromycin | Tablet | Slow-release (b) |
| Klor-Con | potassium | Tablet | Extended-release (b) |
| Klor-Con M | potassium | Table | Extended-release (b,h) |
| Kombiglyze XR | combination | Tablet | Extended-release |
| | | | Note: tablet matrix may remain in stool |
| K-Dur | potassium | Tablet | Extended-release |
| K-Lyte | potassium | Tablet | Effervescent tablet (f) |
| K-Lyte CL | potassium | Tablet | Effervescent tablet (f) |
| K-Lyte DS | potassium | Tablet | Effervescent tablet (f) |
| K-Tab | potassium | Tablet | Extended-release (b) |
| La MIC tal XR | lamo TRI gine | Tablet | Extended-release |
| Lescol XL | fluvastatin | Tablet | Extended-release |
| Letairis | ambrisentan | Tablet | Slow-release |
| Levbid | hyoscyamine | Tablet | Extended-release (h) |
| Levsinex | hyoscyamine | Capsule | Extended-release |
| Lialda | mesalamine | Tablet | Delayed-release |
| Liquibid | combination | Tablet | Extended-release (h) |
| Lithobid | lithium | Tablet | Extended-release |
| Lodrane 24 | brompheniramine | Capsule | Extended-release |
| Lodrane 24D | combination | Capsule | Extended-release |
| LoHist 12 Hour | brompheniramine | Tablet | Extended-release |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|-----------------------------------|-----------------------------------|-----------------|--|
| Lovaza | combination | Capsule | Note: contents of capsule may erode walls of |
| | | | styrofoam or plastic materials |
| Luvox CR | fluvoxa MINE | Capsule | Extended-release |
| Maxifed DM | combination | Tablet | Slow-release (h) |
| Maxifed DMX | combination | Tablet | Slow-release (h) |
| Maxiphen DM | combination | Tablet | Slow-release (h) |
| Mestinon ER | pyridostigmine | Tablet | Extended-release (b) |
| Metadate ER | methylphenidate | Tablet | Extended-release |
| Metadate CD | methylphenidate | Capsule | Extended-release (a) |
| Methylin ER | methylphenidate | Tablet | Extended-release |
| Metoprolol ER | | Tablet | Extended-release |
| Micro K Extencaps | potassium chloride | Capsule | Extended-release (a,b) |
| Mirapex ER | pramipexole | Tablet | Extended-release |
| Moxatag | amoxicillin | Tablet | Extended-release |
| Morphine sulfate extended-release | | Tablet | Extended-release |
| Motrin | ibuprofen | Tablet | Taste (e) |
| MS Contin | morphine | Tablet | Extended-release (b) |
| Mucinex | guia FEN esin | Tablet | Slow-release |
| Mucinex DM | combination | Tablet | Slow-release |
| Myfortic | mycophenolate | Tablet | Delayed-release |
| Namenda XR | memantine | Capsule | Extended-release (a) |
| Naprelan | naproxen | Tablet | Extended-release |
| NexIUM | esomeprazole | Capsule | Delayed-release (a) |
| Niaspan | nicotinic acid | Tablet | Extended-release |
| Nicotinic Acid | | Capsule; Tablet | Slow-release (h) |
| Nifediac CC | NIFEDipine | Tablet | Extended-release |
| Nifedical XL | NIFED ipine | Tablet | Extended-release |
| Nitrostat | nitroglycerin | Tablet | Sublingual route (g) |
| Norflex ER | orphenadrine | Tablet | Extended-release |
| Norpace CR | disopyramide | Capsule | Extended-release form within a special capsule |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|---------------------------|-----------------------------------|----------------|--|
| Norvir | ritonavir | Tablet | Note: crushing tablets has resulted in decreased |
| | | | bioavailability of drug (b) |
| Nucynta ER | tapentadone | Tablet | Extended-release |
| | | | Note: toxic dose may occur if tablet is split or |
| | | | crushed |
| Oleptro | tra ZOD one | Tablet | Extended-release |
| Opana ER | oxymorphone | Tablet | Extended-release |
| Oracea | doxycycline | Capsule | Delayed-release |
| Oramorph SR | morphine | Tablet | Extended-release (b) |
| Orphenadrine citrate ER | | Tablet | Extended-release |
| Oxy CONTIN | oxy CODONE | Tablet | Extended-release |
| | | | Note: tablet disruption may cause a potentially |
| | | | fatal overdose of oxyCODONE |
| Oxymorphone ER | | Tablet | Extended-release |
| Pancrease Delayed-Release | pancrealipase | Capsule | Extended-release |
| Pancrease MT | pancrealipase | Capsule | Enteric-coated (a) |
| Pancrecarb | pancrealipase | Capsule | Enteric-coated (a) |
| Pancrelipase | | Capsule | Enteric-coated (a) |
| Paxil CR | PARoxetine | Tablet | Extended-release |
| Pentasa | mesalamine | Capsule | Slow-release |
| Pradaxa | dabigatran | Capsule | Note: bioavailability increases by 75% when the |
| | | | pellets are taken without the capsule shell |
| Pre-Hist-D | combination | Tablet | Extended-release (h) |
| Plendil | felodipide | Tablet | Extended-release |
| Prevacid | lansoprazole | Capsule | Delayed-release |
| Prevacid SoluTab | lansoprazole | Tablet | Note: Orally disintegrating do not swallow; |
| | | | dissolve in water only and dispense |
| | | | via dosing syringe or NG tube |
| Prevacid Suspension | lansoprazole | Suspension | Slow-release |
| | | | Note: contains enteric-coated granules; mix with |
| | | | water only; not for NG use |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|----------------------|-----------------------------------|----------------|---|
| Pri LOSEC | omeprazole | Capsule | Delayed-release (a) |
| Pri LOSEC OTC | omeprazole | Tablet | Delayed-release |
| Pristiq | desvenlafaxine | Tablet | Extended-release |
| Procardia XL | NIFED ipine | Tablet | Extended-release |
| Propecia | finasteride | Tablet | Note: women who are, or may become, pregnant, should not handle crushed or broken tablets |
| Proquin XR | ciprofloxicin | Tablet | Extended-release |
| Proscar | finasteride | Tablet | Note: women who are, or may become, pregnant, should not handle crushed or broken tablets |
| Protonix | pantoprazole | Tablet | Slow-release |
| PROzac Weekly | FLU oxetine | Tablet | Enteric-coated |
| Qdall | combination | Capsule | Extended-release |
| Qdall AR | combination | Capsule | Extended-release |
| Ranexa | ranolazine | Tablet | Slow-release |
| Rapamune | sirolimus | Tablet | Note: pharmacokinetic NanoCrystal technology may be affected (b) |
| Rayos | predni SONE | Tablet | Delayed-release Note: release is dependent upon intact coating |
| Razadyne ER | galantamine | Capsule | Extended-release |
| Renagel | sevelamer | Tablet | Note: tablets expand in liquid if broken or crushed |
| Renvela | sevelamer carbonate | Tablet | Note: tablets expand in liquid if broken or crushed (b) |
| Requip XL | r OPINIR ole | Tablet | Extended-release |
| Rescon | combination | Tablet | Slow-release (h) |
| Rescon JR | combination | Tablet | Slow-release (h) |
| Rescon MX | combination | Tablet | Slow-release (h) |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|---------------------|-----------------------------------|----------------|--|
| Respahist | combination | Capsule | Extended-release (a) |
| Respaire SR | combination | Capsule | Extended-release |
| Revlimid | lenalidomide | Capsule | Note: Teratogenic potential; healthcare workers should avoid contact with capsule contents/body fluids |
| Ritalin LA | methylphenidate | Capsule | Extended-release (a) |
| Ritalin SR | methylphenidate | Tablet | Extended-release |
| R-Tanna | combination | Tablet | Slow-release |
| Rythmol SR | propafenone | Capsule | Extended-release |
| Ryzolt | tra MAD ol | Tablet | Extended-release |
| | | | Note: crushing may cause overdose |
| Sensipar | cinacalcet | Tablet | Note: tablets are not scored and cutting |
| | | | may cause variable dosage accuracy |
| SERO quel XR | QUE tiapine | Tablet | Extended-release |
| Sinemet CR | levo/carbidopa | Tablet | Extended-release (h) |
| Sinuvent PE | combination | Tablet | Extended-release (h) |
| Slo-Niacin | nicotinic acid | Tablet | Slow-release (h) |
| Solodyn | minocycline | Tablet | Extended-release |
| Somnote | chloral hydrate | Capsule | Liquid filled |
| Sprycel | dasatinib | Tablet | Film-coated Note: active ingredients are surrounded by a wax matrix to prevent healthcare exposure; women who are, or may become, pregnant, should not handle crushed or broken tablets |
| Strattera | atomoxetine | Capsule | Note: capsule contents can cause ocular irritation |
| Sudafed 12 hour | combination | Capsule | Extended-release (b) |
| Sudafed 24 hour | combination | Capsule | Extended-release (b) |
| Sular | nisoldipine | Tablet | Extended-release |
| Symax Duotab | hyoscyamine | Tablet | Controlled-release |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|---------------------|-----------------------------------|-----------------|---|
| Symax SR | hyoscyamine | Tablet | Extended-release |
| Tasigna | nilotinib | Capsule | Note: disruption of capsule may yield high blood |
| | | | levels causing enhanced toxicity |
| Taztia XT | diltiazem | Capsule | Extended-release (a) |
| TEG retol-XR | car BAM azepine | Tablet | Extended-release (b) |
| Temodar | temozolomide | Capsule | Note : accidentally opened or damaged capsules require rigorous precautions to avoid inhalation or contact with the skin or mucous membranes (i) |
| Tessalon Perles | benzonatate | Capsule | Note: swallow whole; local anesthesia of the oral mucosa; choking could occur |
| Theo-24 | theophylline | Capsule | Extended-release Note: contains beads that dissolve throughout the GI tract |
| Theochron | theophylline | Tablet | Extended-release |
| Tiazac | diltiazem | Capsule | Extended-release (a) |
| Topamax | topiramate | Tablet; Capsule | Taste; Taste (a) |
| Toprol XL | metoprolol | Tablet | Extended-release (h) |
| Touro CC-LD | combination | Tablet | Extended-release (h) |
| Touro LA-LD | combination | Tablet | Extended-release (h) |
| Toviaz | fesoterodine | Tablet | Extended-release |
| Tracleer | bosentan | Tablet | Note : women who are, or may become, pregnant, should not handle crushed or broken tablets |
| TRENtal | pentoxifylline | Tablet | Extended-release |
| Treximet | combination | Tablet | Note: unique drug matrix enhances rapid drug absorption |
| Trilpix | fenofibric | Capsule | Extended-release |
| Tylenol Arthritis | acetaminophen | Tablet | Controlled-release |



| Drug Product | Active Ingredient(s) ² | Dosage Form(s) | Reasons/Comments ³ |
|-------------------|-----------------------------------|----------------|--|
| Ultram ER | tra MAD ol | Tablet | Extended-release |
| | | | Note: tablet disruption may cause a potentially |
| | | | fatal overdose of drug |
| Ultrase | pancrealipase | Capsule | Enteric-coated |
| Uniphyl | theophylline | Tablet | Slow-release |
| Urocit-K | potassium citrate | Tablet | Wax-coated; prevents upper GI release |
| Uroxatral | alfuzosin | Tablet | Extended-release |
| Valcyte | val GANC iclovir | Tablet | Teratogenic and irritant potential (i,b) |
| Verapamil SR | | Tablet | Extended-release (h) |
| Verelan | verapamil | Capsule | Sustained-release (a) |
| Verelan PM | verapamil | Capsule | Extended-release (a) |
| Videx EC | didanosine | Capsule | Delayed-release |
| Vimovo | naproxen / esomeprazole | Tablet` | Delayed-release |
| Viramune XR | nevirapine | Tablet | Extended-release (b) |
| Voltaren XR | diclofenac | Tablet | Extended-release |
| VoSpire ER | albuterol | Tablet | Extended-release |
| Votrient | pazopanib | Tablet | Note: crushing significantly increases the AUC and |
| | | | Tmax; crushed or broken tablets may cause |
| | | | dangerous skin problems |
| Wellbutrin SR, XL | bu PROP ion | Tablet | Extended-release |
| | | | |
| Xanax XR | ALPRAZ olam | Tablet | Extended-release |
| Zegerid OTC | omeprazole/ NaHCO3 | Capsule | Delayed-release (b) |
| Zenpep | pancrealipase | Capsule | Delayed-release (a) |
| Zolinza | vorinostat | Capsule | Note: irritant; avoid contact with skin or mucous |
| | | | membranes; avoid contact with crushed or broken |
| | | | tablets |
| Zortress | everolimus | Tablet | Note: crushed powder may cause |
| | | | dangerous effects to mucus membranes |
| Zyban | bu PROP ion | Tablet | Slow-release |
| Zyflo CR | zileuton | Tablet | Extended-release |



- (a) Capsule may be opened and the contents taken without crushing or chewing; soft food such as applesauce or pudding may facilitate administration; contents may generally be administered via nasogastric tube using an appropriate fluid provided entire contents are washed down the tube.
- (b) Liquid dosage forms of the product are available; however, dose, frequency of administration and manufacturers may differ from that of the solid dosage form.
- (C) Antacids and/or milk may prematurely dissolve the coating of the tablet.
- (d) Capsule may be opened and the liquid contents removed for administration.
- (e) The taste of this product form would likely be unacceptable to the patient; administration via nasogastric tube should be acceptable.
- (f) Effervescent tablets must be dissolved in the amount of diluent recommended by the manufacturer.
- (g) Tablets are made to disintegrate under the tongue.
- (h) Tablet is scored and may be broken in half without affecting release characteristics.
- (i) Skin contact may enhance tumor production; avoid direct contact.

Disclaimer: This listing is not meant to represent all products, either by generic or trade name. The author encourages manufacturers, pharmacists, nurses, and other health professionals to notify him of any changes or updates.

1. Correspondence regarding this list may be addressed to:

John F. Mitchell, PharmD, FASHP Email: rxmitchell@att.net

- 2. The generic name is provided merely as a reference point and is only listed for single ingredient medications; it should not be assumed that drugs with the same generic are equivalent to the specific brand name listed relative to crushing or chewing. If questions arise, please check with your pharmacist.
- 3. Two official USP terms are used to designate special-release medication forms: "extended release" and "delayed release". Others such as "sustained release", "controlled release", etc. are commonly used on package labeling. The term "Slow-release" is being used here to signify all such drugs with a special-release mechanism.
- © **Copyright 2011, 2012**. No part of this list may be reproduced in any format without the expressed permission of the author.

